

# TRANSMISSION ELECTRON MICROSCOPIC STUDIES FOR THE IDENTIFICATION OF BACTERIAL ISOLATE FROM RHIZOSPHERE OF BRINJAL (*SOLANUM MELONGENA* L.)

JENIFER LOLITA. C<sup>1</sup> & KESHAMMA E<sup>2\*</sup>

<sup>1</sup>Department of Botany, Maharani Cluster University, Palace Road, Bengaluru, Karnataka, India

<sup>2</sup>Department of Biochemistry, Maharani Cluster University, Palace Road, Bengaluru, Karnataka, India

## ABSTRACT

**Background:** Rhizosphere, phylloplane and caulosphere is the region where a complex community of microbes, mainly bacteria and fungi are present. The microbe plant interaction in these regions can be beneficial, neutral, variable, or deleterious for plant growth. The bacteria that exert beneficial effects on plant development are termed plant growth promoting bacteria. **Objectives:** To identify the bacteria isolated from rhizosphere of brinjal (*Solanum melongena* L.) by transmission electron microscope (TEM). **Materials and Methods:** The pure cultures of bacterial isolate from brinjal (*Solanum melongena* L.) were used to identify the bacteria. Identification of bacteria was done by transmission electron microscope using reference strain viz., *Bacillus polymyxa* strain 10401 obtained from France. **Results:** The BBI revealed rod shaped sporulating bacteria with single polar flagella as observed under transmission electron microscope. The bacterial isolate which showed nitrogen fixing and phosphate solubilizing properties was identified as *Bacillus polymyxa* belonging to the class Eubacteriales, family Bacillaceae and genera Bacillus. **Conclusion:** Transmission electron microscope studies revealed that the BBI was a rod shaped bacteria with a single polar flagellum. The bacterial cell measured 2.57  $\mu\text{m}$  (2570 nm) in length and 0.8570  $\mu\text{m}$  (857 nm) in breadth. Furthermore, TEM studies of the sporulating culture revealed the presence of a large central spore, confirming light microscopic observations. The single large spore occupied the entire inner space of the bacterial cell.

**KEYWORDS:** Brinjal Bacterial isolate (BBI), Transmission Electron Microscopy, Rod Shaped Bacteria & *Bacillus Polymyxa*

**Received:** Oct 12, 2021; **Accepted:** Nov 02, 2021; **Published:** Nov 13, 2021; **Paper Id.:** IJBTRDEC20216

## INTRODUCTION

Brinjal (*Solanum melongena* L.) is a member of the family Solanaceae and a native of India. It is an important vegetable crop of south India. The fruit is rich in vitamin A and vitamin C and is employed in the Ayurveda system. In Karnataka, brinjal is cultivated in 16,602 hectares of land and has an average yield of 30-35 tons per hectare. The crop needs 369 kg of urea and 80 kg of phosphatic fertilizer per hectare. Although symbiotic nitrogen fixation especially legume-rhizobium system has been proved to be the best form of biological nitrogen fixation, associative nitrogen fixation cannot be ignored. Nitrogen fixation on the rhizoplane, phylloplane and stem have been attributed to the presence of diazotrophic bacteria associated with the roots, stem and leaves of plants. [Dobereiner 1980]

There are many reports relating to the characterisation and identification of the nitrogen fixing bacteria associated with a wide variety of grasses and cereal crops. [Dobereiner 1976, Tarrand 1978, van Berkum,

1979, 1980, Silva et al., 1981; Levanony et al. 1987, Rai and Guar, 1988] Hill, 1980; Hill et al., 1983 isolated and characterized rhizosphere bacteria of sweet potato. [Hill et al., 1983] Levanony et al. (1987) identified *Azospirillum brasiliense* in cereal roots using ELISA. Rai and Guar, (1988) characterized *Azotobacter* associated with roots of wheat. Lalande et al. (1989) identified rhizobacteria associated with maize as *Azospirillum*. Lalande R (1989) identified *Azospirillum brasiliense* on the surface and endosphere of wheat roots by immunogold labelling. Brand et al. (1991) isolated a root colonizing rhizobacteria, which was characterized as *Pseudomonas*. McInroy and Kloepper (1991) identified endophytic bacteria of maize and cotton.

Penot et al., 1991 and McInroy and Kloepper, 1991, characterized *Azospirillum* associated with maize cultivated in France, using biochemical tests. Agarwala-Dutt et al. (1991) isolated associated bacteria from the interiors of many graminaceous plants, many of which were identified as *Azospirillum*. Lukin (1990) observed spatial distribution of associated bacteria identified as *Azospirillum brasiliense* the rhizosphere of barley plants. Holguin et al. (1992) isolated and identified rhizobacteria associated with mangrove trees as *Staphylococcus*. Fages and Mulard (1986) identified the isolated bacteria from the rhizosphere of sunflower as *Azospirillum*.

With this background, the present study was designed with the main purpose to identify the brinjal bacterial isolate (BBI) from the rhizosphere of brinjal through transmission electron microscopic (TEM) studies.

## MATERIALS AND METHODS

The isolated BBI was subcultured in solid and liquid nitrogen free liquid Burk's media. The pure cultures of BBI were used to characterize the bacteria. Identification of the dominantly associated bacterial isolate of brinjal (*Solanum melongena* L.) was done comparing with a reference strain viz., *Bacillus polymyxa* strain 10401 obtained from France.

### To Study the Vegetative Cells of BBI

A 24hr pure culture was diluted in 10 ml of sterile distilled water. After thorough shaking, a drop of this suspension was mounted on a copper grid. A drop of the mixture containing bovine serum albumin (0.1%) and PTA with BSA (Buffered phosphotungstate with bovine serum albumin in the ratio of 1:2) was placed on the support film side of the copper grid. After 30 seconds excess fluid was drained off from the grid by touching its edge to the filter paper leaving a very thin film of liquid. The grid was stained with uranyl acetate before drying. The preparation was examined by transmission electron microscope. Several morphological observations like size, shape, presence of flagella was recorded.

### To Study the Sporulating Cells of BBI

A 48hr pure culture of BBI in sporulating media (Brown ME 1968) incubated at 37°C (pH 6.7) was used for the study. The culture was harvested and suspended in physiological saline. One drop of this suspension in four parts of PTA with BSA was mounted on copper grids. After 30 seconds excess fluid was drained off from the grid and stained with uranyl acetate before drying. The preparation was observed under an electron microscope.


## RESULTS

### To Study the Vegetative Cells of BBI

Transmission electron microscope studies confirmed that the BBI was a rod shaped bacteria with a single polar flagellum (Plate-1). The bacterial cell measured 2.57  $\mu\text{m}$  (2570nm) in length and 0.8570  $\mu\text{m}$  (857nm) in breadth.

### To Study the Sporulating Cells of BBI

TEM studies of the sporulating culture revealed the presence of a large central spore, confirming light microscopic observations (Plate-2). The single large spore occupied the entire inner space of the bacterial cell.



**Plate-2: Single cell of BBI Observed under Transmission Electron Microscope showing the Presence of Polar Flagella (1000x)**



**Plate-2: Sporulating cell of BBI Observed under Transmission Electron Microscopy (1400x)**

## DISCUSSIONS

The bacterial isolate which showed nitrogen fixing and phosphate solubilizing properties was identified as *Bacillus polymyxa* belonging to the class Eubacteriales, family Bacillaceae and genera Bacillus (Bergey's Manual of determinative bacteriology 8th edition), which includes another phosphate solubiliser viz., *Bacillus macerans*. *Bacillus polymyxa* appeared to be rod shaped with single polar flagellum with spore observed under a transmission electron microscope. Growth promoting properties of *Bacillus polymyxa* have been reported by Holl and Chanway (1992) in pine seedlings. These studies further support the present finding of growth promotion by *Bacillus polymyxa* isolated from the roots of brinjal (*Solanum melongena* L.). The single spore covered the entire space of the cell. The vegetative cell measured  $2570\mu\text{m} \times 0.8570\mu\text{m}$  and had single long polar flagella that helped in movement and infibrillar attachment. These bacteria are gram +ve, Shawky and Sawaby (1991), Silva et al (1981) and Umali-Gracia (1981) did similar identification of the isolated bacteria using electron microscopy. Shawky and Sawaby (1991) identified the isolate as *Azotobacter* using transmission electron microscopy based on external morphology, sporulation and germination. Levanony et al (1987) used ELISA for specific identification of the rhizobacteria isolated from cereal roots, which was characterized as *Azospirillum*. Growth promoting properties of *Bacillus polymyxa* have been reported by Holl and Chanway (1992), in pine seedlings. These studies further support the present finding of growth promotion by *Bacillus polymyxa* isolated from the roots of brinjal (*Solanum melongena* L.).

## CONCLUSIONS

Transmission electron microscope studies revealed that the BBI was a rod shaped bacteria with a single polar flagellum. The bacterial cell measured 2.57  $\mu\text{m}$  (2570 nm) in length and 0.8570  $\mu\text{m}$  (857 nm) in breadth. Furthermore, TEM studies of the sporulating culture revealed the presence of a large central spore, confirming light microscopic observations. The single large spore occupied the entire inner space of the bacterial cell.

## REFERENCES

1. Dobereiner J. Forage grasses and grain crops. *Methods for evaluating biological nitrogen fixation*. 1980:535-6.
2. Dobereiner J, Marriel IE, Nery M. Ecological distribution of *Spirillum lipoferum* Beijerinck. *Canadian Journal of Microbiology*. 1976;22(10):1464-73.
3. Tarrand JJ, Krieg NR, Döbereiner J. A taxonomic study of the *Spirillum lipoferum* group, with descriptions of a new genus, *Azospirillum* gen. nov. and two species, *Azospirillum lipoferum* (Beijerinck) comb. nov. and *Azospirillum brasilense* sp. nov. *Canadian Journal of Microbiology*. 1978;24(8):967-80.
4. van Berkum P, Sloger C. Immediate acetylene reduction by excised grass roots not previously preincubated at low oxygen tensions. *Plant Physiology*. 1979;64(5):739-43.
5. Van Berkum P, Bohlool B. Evaluation of nitrogen fixation by bacteria in association with roots of tropical grasses. *Microbiological reviews*. 1980; 44(3):491-517.
6. Silva DM, Ruschel AP, Matsui E, Lima Nogueira ND, Vose PB. Determination of the activity of N<sub>2</sub>-fixing bacteria in sugarcane roots and bean nodules using tritiated acetylene reduction technique and electron microautoradiography. In *Associative Nitrogen fixation* Eds. 1981:145-152.
7. Levanony H, Bashan Y, Kahana ZE. Enzyme-linked immunosorbent assay for specific identification and enumeration of *Azospirillum brasilense* Cd. in cereal roots. *Applied and Environmental Microbiology*. 1987;53(2):358-64.
8. Rai SN, Gaur AC. Characterization of *Azotobacter* spp. and effect of *Azotobacter* and *Azospirillum* as inoculant on the yield and N-uptake of wheat crop. *Plant and Soil*. 1988;109(1):131-4.
9. Hill WA, Bacon-Hill P, Crossman SM, Stevens C. Characterization of N<sub>2</sub>-fixing bacteria associated with sweet potato roots. *Canadian Journal of Microbiology*. 1983; 29(8):860-2.
10. Lalonde, R., Bissonnette, N., Coutlee, D. and Antoun, H. (1989). *Plant and Soil* (Netherlands). 1989;117 (2): 207-218.
11. Brand I, Lugtenberg BJ, Glandorf DC, Bakker PA, Schippers B, De Weger LA. Isolation and characterization of a superior potato root-colonizing *Pseudomonas* strain. *Bulletin OILB SROP* (France). 1991.
12. McInroy JA, Kloepper JW. Analysis of population densities and identification of endophyte bacteria of maize and cotton in the field. *Bulletin OILB SROP* (France). *OILBSIRP*. 1991:328- 331.
13. Penot I, Berges N, Guinguene C, Fages J. Characterization of *Azospirillum* associated with maize (*Zea mays*) in France, using biochemical tests and plasmid profiles. *Canadian Journal of Microbiology*. 1992;38(8):798-803.
14. Agarwala-Dutt R, Tilak KV, Rana JP. Isolation of *Azospirillum* from the interior of various parts of some graminaceous plants. *Zentralblatt für Mikrobiologie*. 1991; 146(3):217-9.
15. Lukin S, Kozhevnikov P, Zviagintsev D. Spatial distribution of *Azospirillum brasilense* cells in the rhizosphere of barley plants. *Mikrobiologiya* (Moskva, 1932). 1990(6):1090-3.

16. Holguin G, Guzman MA, Bashan Y. Two new nitrogen-fixing bacteria from the rhizosphere of mangrove trees: Their isolation, identification and in vitro interaction with rhizosphere *Staphylococcus* sp. *FEMS Microbiology Letters*. 1992;101(3):207-16.
17. Fages J, Mulard D. Isolation of rhizosphere bacteria and their effect on *Zea mays* in pots [*Azospirillum lipoferum*, *Enterobacter cloacae*, *Pseudomonas diminuta*]. *Agronomie (France)*. 1988; 1108(4):309-314.
18. Rao NS. Nitrogen-fixing bacteria associated with plantation and orchard plants. *Canadian Journal of Microbiology*. 1983;29(8):863-6.
19. Brown ME. Production of plant growth substances by *Azotobacter chroococcum*. *Microbiology*. 1968 Aug 1;53(1):135-44.
20. Krieg NR, Holt JG. *Bergey's manual of systematic bacteriology*. Yi Hsien Publishing Co. 1984.
21. Holl FB, Chanway CP. Rhizosphere colonization and seedling growth promotion of lodgepole pine by *Bacillus polymyxa*. *Canadian Journal of Microbiology*. 1992;38(4):303-8.
22. Shawky BT, Sawaby A. Transmission electron microscopic studies on the external morphology of *Azotobacter* during encystment and germination. *Zentralblatt für Mikrobiologie*. 1991;146(7-8):495-507.
23. Umali-Garcia M, Hubbell DH, Gaskins MH, Dazzo FB. Adsorption and mode of entry of *Azospirillum brasilense* to grass roots [Nitrogen fixation]. 1981.
24. Afzal, A. Alia, et al. "Phosphate solubilizing bacteria associated with vegetables roots in different ecologies." *Pak. J. Bot* 45.SI (2013): 535-544.
25. Naraghi, Laleh, et al. "Increasing the Stability of Antagonistic Fungus, *Talaromyces Flavus*, Using Some Organic and Mineral Compounds." *International Journal of Agricultural Science and Research (IJASR)* 8.1 (2018): 181-190.
26. Sathya, S., S. Lakshmi, and S. Nakkeeran. "Combined effect of biopriming and polymer coating against chilli damping off." *International Journal of Agricultural Science and Research*, 6 (3), 45 54 (2016).
27. Kumar, M. Sunil, and R. Sarada Jayalakshmi Devi. "A Review on Colletotrichum Blight of Chickpea in Andhra Pradesh." *International Journal of Agricultural Science and Research (IJASR)* 6 (2016): 145-158.

